

KCL urban model description, evaluation and outputs

Model summary

The KCLurban model used a kernel modelling technique, based upon the ADMS^{1,2}, to describe the initial dispersion from each emissions source. The contribution from each source has been summed onto a fixed 20m x 20m grid across London assuming that one can calculate the contribution of any source to total air pollution concentrations by applying each kernel and adjusting for the source strength. The kernels were produced using an emissions source of unity, either 1 g s^{-1} (point and jet sources), $1 \text{ g m}^{-3} \text{ s}^{-1}$ (volume sources) or a $1 \text{ g km}^{-1} \text{ s}^{-1}$ (road sources) and hourly meteorological measurements from the UK Meteorological Office site at Heathrow. Data from the Heathrow site is recorded at a height of 10 metres and includes measurements of temperature, wind speed, wind direction, precipitation, relative humidity and cloud cover. The KCLurban model used emissions from the London Atmospheric Emissions Inventory (LAEI)³ and for road traffic emissions using King's well established emissions modelling methods, combined with emissions factors from the UK specific roadside measurements of Carslaw⁴ and for non-exhaust emissions, based upon the work of Harrison⁵.

Treatment of sources in KCLurban

To predict air quality concentrations in London the KCLurban model sums together three source categories. First, for rural sources outside the model domain, use was made of rural NO_x , PM_{10} , $\text{PM}_{2.5}$ and monthly OX ($\text{NO}_2 + \text{O}_3$), measurements taken from a combination of Harwell, Rochester Stoke and Maidstone monitoring sites, part of the UK AURN network (web link.....). Second, within the model domain, but greater than 500m from a receptor location, London sources were represented as shallow volumes of 1x1 km horizontal dimension and 2m vertically for road traffic, and 50m vertically for other sources. Because emissions from biomass burning are not well represented in the London emissions inventory an additional $1.05 \mu\text{g m}^{-3}$ of PM was also included, based upon the work of Fuller⁴. Third, for those sources within 500m of a receptor location, a detailed treatment of road/rail/aircraft and gas combustion sources was used, each of which is described in sections 1.3 to 1.6.

Sources within the KCLurban model include: road transport (exhaust and non-exhaust), large regulated industrial processes, small regulated industrial processes, large boiler plant, gas heating (domestic and industrial-commercial), oil combustion sources (domestic and commercial), coal combustion sources (domestic and commercial), agricultural and natural sources, rail, ships, airports and others (sewage plant etc). In modelling the emissions from large industrial processes use was made of emissions data and stack conditions (height, temperature, volume flow rate) for each source.

Representing road sources

Within 500m of a road, where strong concentration gradients exist, a highly detailed treatment of road sources is required. In KCLurban we modelled road emissions as a series of road links 10 m long and based on geographically accurate Ordnance Survey road map data. To reflect the important separation of two or more lanes of traffic, on large roads/motorways, each side of the road was modelled separately. Six road categories (and associated kernels) were used in London, including open roads (motorway), typical roads (average urban roads surrounded by low rise buildings) and 4 types of street canyon (classified by their orientation: north-south, east-west, southwest-northeast and southeast-northwest). The “typical roads” had a road width of 20m and a building height of 10m and “street canyons” had a width of 30m and a building height of 25m. The “street canyon” width and height details were manually sampled from the 3D building model in London (<http://www.casa.ucl.ac.uk/news/newsStory.asp?ID=80>). There are approximately 2.25 million 10m road sources in the KCLurban model.

Representing railway sources

Railway sources were treated in much the same way as for roads, i.e. by using the rail network emissions broken into 10m sections. Whilst trains in London are predominantly electric, the line running west from Paddington station and the line running north from St Pancras station includes many diesel trains and represents a significant source of NO_x and PM emissions. For diesel trains, the emissions release height was taken to be 5m and the NO₂/NO_x exhaust ratio was assumed to be similar to large HGVs at 13%.

Representing gas combustion sources

Gas combustion is a very important source of NO_x in London³, and in the KCLurban model a detailed representation of the height of release, spatial distribution of gas sources and temporal change in emissions throughout the year has been included. Through analysis of the 3D model of buildings in London (<http://www.casa.ucl.ac.uk/news/newsStory.asp?ID=80>) the height of release from gas sources was varied from 1m (domestic housing), through 30m for small commercial premises to 75m for large commercial office buildings. Gas heating sources were represented spatially by points located at 50m intervals throughout the minor road network and set back from the road by 20m, assuming that the majority of buildings follow the road network. Model kernels representing the varied release conditions from these sources were then applied to domestic and commercial gas emissions summarised in 1 x 1km grids across London. Finally, a detailed treatment of the emissions variation by hour of the day and month of the year was included in the model and was based upon UK gas use statistics (http://www.decc.gov.uk/en/content/cms/statistics/energy_stats/source/gas/gas.aspx).

Representing Heathrow airport

At Heathrow airport, emissions from aircraft during approach, landing, taxi out, taxi in, hold, take off, initial climb and climb out were represented using the same method as for roads, i.e. as individual sources 10m apart. Auxiliary Power Unit (APU) emissions and engine testing on the airport were

represented as stationary point sources. Other sources such as heating plant, public and staff car parks, car rental, taxis queues and fire training ground emissions were represented horizontally as volume sources of 1 x 1 km, 50m high, and for airside vehicles 2m high.

Takeoff is the aircraft mode that provides one of the largest contributions to ground-level NO_x and PM concentrations. During takeoff each accelerating aircraft engine was represented by horizontal stationary jet sources at 10 m intervals along the runway, from the start of the ground-roll to where the aircraft leaves the ground. The effect of accelerating aircraft was included to reflect the maximum emission close to the start of the ground-roll, thereby reproducing the steep emission gradient along the runway.

Of importance in estimating ground level concentrations from Jet plumes is their rapid change in temperature and velocity over short distances, as well as the effect of aircraft speed. By comparing modelled and measured data close to Heathrow airport we showed that buoyant jet plumes could be accurately modelled by varying the jet release temperature assumptions. The hourly variation of aircraft emissions was reproduced using aircraft movements made available by the UK Civil Aviation Authority. Jet velocities were varied for the different aircraft operational settings of take off, approach and taxiing and were assumed to be 85 %, 30 % and 7 % of full thrust, respectively. Finally, account was taken of the rapidly reducing effect of aircraft emissions to ground level concentrations at different aircraft heights. The aircraft NO₂/NO_x emissions ratio was assumed to be 11%.

Predicting monthly mean NO₂ and O₃ concentrations

The method for converting NO_x to NO₂ used the well established relationships of Carslaw et al⁶ and was based upon an analysis of measurements at both background and roadside sites, included the influence of NO₂ emitted directly from the vehicle exhaust. O₃ concentrations were predicted through methods described in Clapp and Jenkins⁷, combined with knowledge of monthly OX entering London, and taken from the Harwell monitoring site, the model's predictions of NO₂ and at roadside the magnitude of primary NO₂ emitted from vehicle exhausts.

KCLurban model evaluation


The KCLurban model is well established for use in public health research (add some refs) and has been submitted to the UK Model Inter-comparison Exercise, run by King's on behalf of DEFRA (http://uk-air.defra.gov.uk/library/reports?report_id=777). However, as part of the Traffic project, the model has been developed to provide monthly average concentrations between 2003 and 2010 and so the following section describes the model's performance, focusing aspects, which are most important to the epidemiology research being undertaken. We therefore describe KCLurban's monthly performance across all sites, using the correlation coefficient *r* and normalised mean bias, its performance across the measurement site types, roadside, urban background and suburban and its seasonal performance.

KCLurban's monthly evaluation across London.

For each of the 96 months between 2003 and 2010, a comparison of observed vs. modelled concentrations has been created and the spearman correlation coefficient (r) and the normalised mean bias (NMB) calculated. The values of r for NO_x and NO_2 vary across a relatively small range giving an average and (standard deviation) of 0.82 (0.049) and 0.84 (0.033). The model performs slightly less well for O_3 concentrations but still results in an average r value of 0.77 (0.075). The NMB for NO_x and NO_2 is centred close to zero at 1% (12%) and 5.5% (13%) with the O_3 predictions having a negative bias of -16% (24%). The values of r for PM_{10} also vary across a relatively small range giving an average and (standard deviation) of 0.72 (0.07) with $\text{PM}_{2.5}$ having a slightly higher value of r of 0.73 and with a higher standard deviation of 0.19. There is small positive NMB for PM_{10} and $\text{PM}_{2.5}$ of 6% (10%) and 4% (9%).

KCLurban's monthly evaluation between site types

As much of the epidemiological analysis undertaken in the Traffic project relies upon the spatial contrast in concentrations by location in London, further analysis of the model performance at roadside, urban background and suburban sites has been undertaken. Roadside locations represent concentrations at approximately 5m from the kerb, urban background sites at locations which are unaffected by single local sources and are within the urban centre and suburban sites are also unaffected by a single local source but on the outskirts of London.






Figure 1 Scatter plots of observed vs. modelled average monthly concentrations for each of the 96 months between 2003 and 2010 by site type: roadside, urban background and suburban

Table 1. Values of the spearman correlation coefficient (r), root mean square error (RMSE) and normalised mean bias (NMB) for observed vs. modelled average monthly concentrations for each of the 96 months between 2003 and 2010 by site type: roadside, urban background and suburban

Site Type	Pollutant	Number of months	Normalised mean bias (NMB)	Root mean square error (RMSE)	Spearman correlation coefficient (r)
Roadside	NO _x	96	-0.02	8.64	0.93
Suburban	NO _x	96	0.17	7.36	0.94
Urban background	NO _x	96	0.20	10.39	0.93
Suburban	NO ₂	96	0.17	3.84	0.91
Urban background	NO ₂	96	0.12	4.17	0.88

Roadside	NO ₂	96	0.06	4.75	0.83
Suburban	PM ₁₀	96	0.06	2.71	0.91
Urban background	PM ₁₀	96	0.07	2.83	0.93
Roadside	PM ₁₀	96	0.06	3.15	0.92
Roadside	PM _{2.5}	87	0.03	1.44	0.94
Urban background	PM _{2.5}	94	0.06	1.87	0.93
Suburban	PM _{2.5}	96	0.06	1.52	0.94
Suburban	O ₃	96	-0.14	3.96	0.92
Urban background	O ₃	96	-0.14	3.85	0.90
Roadside	O ₃	95	-0.12	3.48	0.91

For all of the pollutants the performance of the model (see Figure 1 and Table 1) is good and especially so for NO_x, PM₁₀ and PM_{2.5} ($r > 0.91$), reflecting the contrast between different locations within London and providing robust metrics for subsequent epidemiology analysis. NO₂ ($r > 0.83$) and O₃ ($r > 0.9$) predictions are also reasonably robust and reflect the different London environments. There is some positive bias in the NO_x model at background locations (~18.5%) but very little at roadside (-2%). For NO₂ the NMB is smaller at background sites (~14.5%) and (~6%) at roadsides. Both PM₁₀ and PM_{2.5} metrics have a small positive bias (~6%) across all site types. This suggests that for NO_x the spatial contrast is slightly smaller than the measurements would suggest and that for the PM metrics the spatial contrast should be the most robust. For O₃ there is a consistent negative bias in the predictions across all site types, reflecting the positive NO₂ bias at background locations and the small under prediction of NO_x at roadsides, so the model slightly under predicts O₃ but shows robust spatial contrast for this pollutant.

For some of the months with the highest concentrations the model tends to over predict, with some evidence that this also occurs for NO_x. There are a number of possible reasons for this including the London emissions of NO_x and errors associated with the dispersion model. Also, given that this slight over prediction is evident at suburban locations, the choice of NO_x concentrations coming into London, taken from rural measurements, could also be a source of error. The lowest monthly O₃ predictions have a negative bias and reflect the positive bias in predictions of NO₂ and NO_x.

KCLurban's seasonal performance

As a further insight into the performance of the KCLurban model, we have plotted a time series of monthly average model predictions and observations for all pollutants: NO_x, NO₂, O₃, PM₁₀ and PM_{2.5}, separated by location: roadside, urban background and suburban and by year: 2003 to 2010. For the sake of clarity, we have only included one example for each pollutant, in Figures 2 - 6.




Figure 2 Monthly average measured (triangles) and modelled (circles) NO_x concentrations (ppb) at roadside, urban background and suburban locations for 2003 to 2010

The KCLurban monthly model predictions, summarised in Figure 2, are in good agreement with the observations at roadside sites for all years between 2003 and 2010. However, there are a small number of individual months, when the model exhibits noticeable bias, including during the winter months, October to February, during the first half of 2003 and for urban background and suburban sites during February to May 2004. Comparison of the monthly averaged modelled and measured concentrations of NO₂ follow those of NO_x and for the sake of clarity are not shown, although a positive bias in modelled NO₂ concentrations is also noticeable during the months October to December in 2005 and 2007.

A comparison between PM_{2.5} modelled and measured concentrations show that for only a relatively small number of months, mainly in 2004 and at urban background and suburban sites, is there a notable positive bias in the model results. Care needs to be taken with over interpreting these results as in 2003 and 2004, only 3 and 8 sites existed in London, this being clearly demonstrated by the 2003 roadside results, which were for one site with limited data capture. The results for PM₁₀ have not been shown, but show closer agreement between modelled and measured concentrations than for PM_{2.5}, with no single period of model bias being apparent between 2003 and 2010. This is partially at least a consequence of having greater than 45 PM₁₀ measurement sites in London and also that the limitations of the measurements, in particular loss of semi volatile PM components has been corrected using methods developed by Green⁸.




Figure 3 Monthly average measured (triangles) and modelled (circles) $PM_{2.5}$ concentrations ($\mu g m^{-3}$) at roadside, urban background and suburban locations for 2003 to 2010




Figure 4 Monthly average measured (triangles) and modelled (circles) O_3 concentrations (ppb) at roadside, urban background and suburban locations for 2003 to 2010

The KCLurban model's monthly O_3 results are in reasonable agreement with observations in London although some periods during years 2005 and 2009 exist showing the model having negative bias and in particular in 2005 and 2008.

KCLurban's annual performance

As part of the Traffic project, the KCLurban model has also been used to provide annual average concentrations between 2003 and 2010 and so the following section describes the annual model performance. To understand the annual performance of the KCLurban model predictions, an evaluation process has been undertaken against fixed site measurements from the London Air Quality Network (LAQN) and Defra's Automatic Urban and Rural Network (AURN) covering all the years from 2003 to 2010. This has been based upon our extensive experience of model evaluation, including King's participation in both phases of the UK Model Inter-comparison Exercise, run by King's on behalf of DEFRA (http://uk-air.defra.gov.uk/library/reports?report_id=777).

For NO_x and NO_2 , modelled data has been evaluated against measurement data from monitoring sites with an annual data capture rate of > 75 %. This has resulted in a minimum of 62 site comparisons in 2003 and a maximum of 100 comparisons in 2008 (Table 2, Figure 5 and Figure 6). For both pollutants, a strong linear relationship is observed between the variables and high correlation coefficients were found for both NO_x (varying between 0.79 in 2009 and 0.92 in 2008) and NO_2 (varying between 0.85 in 2009 and 0.93 in 2008). Additionally, for both pollutants the normalised mean gross error (NMGE) is relatively low and for NO_x has a maximum of 23% in 2010 and for NO_2 has a maximum of 16% in 2007. The KCLurban model exhibits little overall bias for NO_2 predictions, with the normalised mean bias (NMB) varying between -8.5 % in 2003 and 6.8 % in 2007, although a small but consistent under prediction of NO_x concentrations is observed: NMB varies between -4.9 % in 2004 and -15.4 % in 2003.

There are fewer measurements of O_3 , with a minimum of 24 monitoring sites in 2003 and a maximum of 37 sites in 2007, all of which have a data capture rate of > 75 % (Table 2, Figure 9). The NMGE varies between 9.2 % in 2007 and 22.1 % in 2010, the NMB varies between -11.5 % in 2003 and 21.7 % in 2010 and the value of r between measured and modelled concentrations varies between 0.79 in 2005 and 0.89 in 2009 and is broadly comparable to that observed for NO_x .

Modelled PM_{10} concentrations have been evaluated against measurements from a minimum of 49 monitoring sites in 2003 and a maximum of 76 sites in 2008, all of which have a data capture rate of > 75 % (Table 2, Figure 7). The correlation coefficient, r , for PM_{10} concentrations indicates a reasonably strong agreement, varying between 0.65 in 2007 and 0.85 in 2005. Overall NMGE in the modelled concentrations is low varying between 7.9 % in 2003 and 13.5 % in 2007, and broadly comparable to that observed for NO_2 . The KCLurban model exhibits little bias in predicting PM_{10} concentrations with NMB varying between -0.87 % in 2003 and 6.8 % in 2006.

There are significantly fewer measurements of $\text{PM}_{2.5}$ than for other pollutants within London. As a result $\text{PM}_{2.5}$ concentrations have been evaluated against measurements from a minimum of only 4 monitoring sites in 2003 and a maximum of 21 sites in 2009, all of which have a data capture rate of > 75 % (Table 2, Figure 8). It is prudent therefore not to over interpret the results for some years. The NMGE varies between 11 % in 2008 and 42 % in 2003 ($n=4$), NMB varies between 0.19 % in 2008 and 42.1 % in 2003 ($n=4$) and the correlation coefficient, r varies between 0.26 in 2010 and 1 in 2004 ($n=5$) is also not as

consistently strong as for NO_x , NO_2 and PM_{10} . It is also important to note that a transition between TEOM based measurements to TEOM-FDMS machine has occurred in a piecemeal way during the period between 2003 and 2010. Whilst the FDMS instruments do not suffer nitrate loss, they are more complex to run and once again add a level of uncertainty to the observations which does not exist for PM_{10} , NO_x , NO_2 or O_3 .

Table 2. Values of the spearman correlation coefficient (r), root mean square error (RMSE), normalised mean gross error (NMGE) and normalised mean bias (NMB) for observed vs. modelled annual average concentrations for each of the years between 2003 and 2010

Year	Pollutant	Number of monitoring sites (n)	Normalised mean bias (NMB)	Normalised mean gross error (NMGE)	Root mean square error (RMSE)	Spearman correlation coefficient (r)
2003	NO _x	62	-0.154	0.21	18	0.88
2004	NO _x	70	-0.049	0.19	17	0.91
2005	NO _x	84	-0.108	0.2	21	0.88
2006	NO _x	85	-0.136	0.21	21	0.86
2007	NO _x	90	-0.069	0.2	18	0.88
2008	NO _x	100	-0.120	0.2	18	0.92
2009	NO _x	96	-0.091	0.23	22	0.79
2010	NO _x	87	-0.132	0.23	22	0.86
2003	NO ₂	62	-0.085	0.13	4.7	0.9
2004	NO ₂	70	0.018	0.12	5.2	0.92
2005	NO ₂	84	-0.015	0.13	6.1	0.91
2006	NO ₂	85	-0.049	0.13	6.1	0.91
2007	NO ₂	90	0.068	0.16	6	0.9
2008	NO ₂	100	-0.011	0.13	5.1	0.93
2009	NO ₂	96	0.031	0.15	5.5	0.85
2010	NO ₂	87	-0.014	0.14	5.8	0.89
2003	PM ₁₀	49	-0.0087	0.079	4	0.71
2004	PM ₁₀	59	0.0604	0.124	4.6	0.76
2005	PM ₁₀	62	0.0391	0.095	3.5	0.85
2006	PM ₁₀	73	0.0068	0.111	5.2	0.73
2007	PM ₁₀	70	0.0441	0.135	5.8	0.65
2008	PM ₁₀	76	-0.0067	0.104	4.1	0.78
2009	PM ₁₀	74	0.0418	0.117	3.7	0.78
2010	PM ₁₀	68	0.0415	0.13	4.2	0.71
2003	PM _{2.5}	4	0.4207	0.42	6.9	0.97
2004	PM _{2.5}	5	0.3474	0.35	5.2	1
2005	PM _{2.5}	9	0.2143	0.26	3.9	0.67
2006	PM _{2.5}	9	0.1533	0.18	2.8	0.84
2007	PM _{2.5}	14	0.0431	0.12	2.1	0.82
2008	PM _{2.5}	15	0.0019	0.11	2	0.74
2009	PM _{2.5}	21	0.104	0.15	2.8	0.59
2010	PM _{2.5}	20	0.0157	0.15	2.8	0.26
2003	O ₃	24	-0.115	0.124	2.7	0.87
2004	O ₃	25	-0.033	0.095	2	0.82
2005	O ₃	29	-0.019	0.095	2	0.79
2006	O ₃	29	-0.047	0.095	2.2	0.81
2007	O ₃	37	0.053	0.092	1.9	0.87
2008	O ₃	36	0.086	0.103	2.3	0.87
2009	O ₃	34	0.138	0.148	2.8	0.89
2010	O ₃	32	0.217	0.221	4	0.85




Figure 5 Scatter plots of annual average modelled vs. measured NO_x concentrations (ppb) for 2003 to 2010




Figure 6 Scatter plots of annual average modelled vs. measured NO_2 concentrations (ppb) for 2003 to 2010




Figure 7 Scatter plots of annual average modelled vs. measured PM_{10} concentrations ($\mu\text{g m}^{-3}$) for 2003 to 2010




Figure 8 Scatter plots of annual average modelled vs. measured $\text{PM}_{2.5}$ concentrations ($\mu\text{g m}^{-3}$) for 2003 to 2010




Figure 9 Scatter plots of annual average modelled vs. measured O₃ concentrations (ppb) for 2003 to 2010

KCLurban's NO₂ model results

The KCLurban model has been used to predict both monthly and annual average concentrations at 20m resolution between 2003 and 2010 as the basis for all of the epidemiological research being undertaken in the Traffic Pollution and Health in London project. This section provides a few examples of the outputs produced by the KCLurban model. Figure 10 illustrates mapped concentrations of NO₂ for the London area up to and including the M25 in the year 2008. The map highlights the importance of road traffic as a source of air pollution. London's major roads are plainly visible and in the majority of cases, the highest concentrations are predicted at, or immediately adjacent to, these major roads. Central London is also highlighted as a problem area. The concentrations in central London are amongst the highest in London and reflect the high density of emission sources. Heathrow airport to the west of the city is also clearly visible, as is the Paddington railway line (running due west from central London).

The KCLurban model outputs have subsequently been manipulated to produce averaged and interpolated concentrations at address and postcode level. Figure 11 shows postcode address centroid annual mean NO₂ concentrations, in the year 2008, interpolated from 20m grid model predictions using a bi-linear interpolation technique.




Figure 10 Annual mean NO₂ concentrations in 2008




Figure 11 Annual mean NO₂ concentrations in 2008 at postcode address centroids

References

1. CERC, 2004, ADMS 4 and Roads user guides. Available at www.cerc.co.uk.
2. CERC. 2006: *ADMS-Roads An Air Quality Management System User Guide*. Cambridge Environmental Research Consultants Ltd, Cambridge. (http://www.cerc.co.uk/environmental-software/assets/data/doc_userguides/CERC_ADMS-Roads%20Extra3.1_User_Guide.pdf). Accessed 03/01/2012
3. Greater London Authority (GLA). 2010. The London Atmospheric Emissions Inventory 2010. <http://data.london.gov.uk/dataset/london-atmospheric-emissions-inventory-2010> Accessed 8/1/2015
4. Carslaw, D.C., Beevers, S.D., Tate, J.E., Westmoreland, E. and M.L. Williams, 2011a. Recent evidence concerning higher NO_x emissions from passenger cars and light duty vehicles. *Atmospheric Environment* 45 (2011) 7053-7063
5. Harrison, R.M. et al (2012) Estimation of the Contributions of Brake Dust, Tire Wear, and Resuspension to Nonexhaust Traffic Particles Derived from Atmospheric Measurements. *ES&T* 46, 6523-6529.
6. Carslaw D.C., Beevers S.D., Fuller G.W., 2001 An empirical approach for the prediction of annual mean nitrogen dioxide concentrations in London, *Atmospheric Environment*, 35(8): 1505-1515.
7. Clapp L.J. and Jenkin M(E. 2001. Analysis of the relationship between ambient levels of O₃, NO₂ and NO as a function of NO_x in the UK. *Atmospheric Environment*, 35 (2001): 6391-6405.
8. Green DC., Fuller GW., Baker T. 2009. Development and validation of the volatile correction model for PM 10 – An empirical method for adjusting TEOM measurements for their loss of volatile particulate matter. *Atmospheric Environment - ATMOS ENVIRON* 01/2009; 43(13):2132-2141.